



Cutaneous lesions in the setting of disseminated *Cryptococcus neoformans* infection

**Authors:**Mithra John¹ Stacey Norsworthy¹ Lauren Richards¹ **Affiliations:**

¹Department of Health,
Internal Medicine,
Helen Joseph Hospital,
Johannesburg, South Africa

Corresponding author:

Mithra John,
mithrajohn10@gmail.com

Dates:

Received: 09 Sept. 2021

Accepted: 21 Sept. 2021

Published: 29 Oct. 2021

How to cite this article:

John M, Norsworthy S,
Richards L. Cutaneous lesions
in the setting of disseminated
Cryptococcus neoformans
infection. S Afr J HIV
Med. 2021;22(1), a1315.
[https://doi.org/10.4102/
sajhivmed.v22i1.1315](https://doi.org/10.4102/sajhivmed.v22i1.1315)

Copyright:

© 2021. The Authors.
Licensee: AOSIS. This work
is licensed under the
Creative Commons
Attribution License.

Read online:

Scan this QR
code with your
smart phone or
mobile device
to read online.

A 38-year-old HIV-positive South African man presented with a 3-week history of headache and worsening widespread cutaneous lesions. He had recently started antiretroviral treatment (ART), was virologically suppressed and had a CD4 T-lymphocyte count of 21 cells/ μ L.

Clinically, the patient had ulcerated, centrally necrotic plaques and papules found mainly on the face and scalp, with lesser involvement of the torso and limbs. The largest lesion was noted to be on his left cheek and measured 4 cm in diameter (Figure 1a). Additionally, the patient displayed signs of meningitis and a right cranial nerve VI palsy on neurological examination.

A diagnosis of disseminated *Cryptococcus neoformans* infection was made based on a positive serum and cerebrospinal fluid *Cryptococcus* antigen and a skin punch biopsy. Histological evaluation of the skin biopsy specimen revealed numerous fungal organisms with a large clear capsule and narrow-based budding on both periodic acid–Schiff and Grocott–Gomori's methenamine silver staining in keeping with *C. neoformans* (Figure 1c and 1d). This was further confirmed on fungal culture of the biopsy specimen.

The patient received amphotericin B deoxycholate and flucytosine for 7 days, as well as daily therapeutic lumbar punctures to relieve raised intracranial pressure. He was then




FIGURE 1: (a) A 4-cm, centrally necrotic plaque on the patient's left cheek with a smaller papule adjacent to it; (b) significant improvement of the same lesions following 5 weeks of therapy; (c) periodic acid–Schiff and (d) Grocott–Gomori stained sections showing numerous fungal organisms with a large clear capsule, in keeping with *Cryptococcus neoformans* (original magnification, 40 \times).

transitioned to oral fluconazole, and ART was reintroduced 5 weeks later. The patient responded well to treatment, with significant improvement to his skin lesions (Figure 1b).

Acknowledgements

The authors thank Dr J. Nel and Dr L.P. Meko for their input in producing this manuscript.

Competing interests

The authors declare that they have no financial or personal relationships that may have inappropriately influenced them in writing this article.

Authors' contributions

M.J. drafted the manuscript. L.R and S.N. reviewed the manuscript.

Ethical considerations

Permission was obtained from the Human Research Ethics Committee (Medical), University of the Witwatersrand, Johannesburg (protocol number M210756).

Funding information

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Data availability

Data sharing is not applicable to this article, as no new data were created or analysed in this study.

Disclaimer

The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of any affiliated agency of the authors.